

L15 ANSWER 41 OF 101 CA COPYRIGHT 2002 ACS  
AN 118:5175 CA  
TI Comparison of Freund's and Ribi adjuvants for inducing antibodies to the  
synthetic antigen (TG)-AL in rabbits  
AU Deeb, Barbara J.; DiGiacomo, Ronald F.; Kunz, Lawrence L.; Stewart, Julie  
L.  
CS Sch. Med., Univ. Washington, Seattle, WA, 98195, USA  
SO J. Immunol. Methods (1992 ), 152(1), 105-13  
CODEN: JIMMBG; ISSN: 0022-1759  
DT Journal  
LA English  
AB Antibody responses and health parameters were compared in rabbits  
immunized with a synthetic polypeptide antigen, [L-Tyr,L-Glu,DL-Ala]-poly-

Applicants: Alexander Gad and Dora Lis  
Serial No.: 09/816,989  
Filed: March 23, 2001  
Exhibit 24

ot

L-lysine [(TG)-AL], in Freund's (FA) or Ribi (RA) adjuvants. Rabbits, 12 wk old, of both sexes, were inoculated with 0.5 mL divided between 2 i.m. sites. Eight received FA and antigen (50 .mu.g); 8 RA and antigen, 8 PBS and antigen; 4 FA and PBS; 4 RA and PBS; and 4 PBS. Identical booster inoculations were made 21 days later, except that incomplete FA was substituted for complete FA. Rabbits were monitored until euthanasia and necropsy 7 wk after the primary inoculation. Sera, obtained weekly, were analyzed for Igs using an enzyme immunoassay. Only rabbits given antigen with adjuvant produced high titered antibodies. Mean optical d. values for IgM were greater the week after the booster in the group given FA. IgG values were similar for both adjuvant/antigen groups the week after the booster, but thereafter decreased in rabbits given RA. Antisera from rabbits given antigen with FA had greater avidity for the antigen than that from rabbits given antigen with RA, however, the difference was not significant. Rabbits inoculated with FA and antigen had high serum creatinine kinase levels the day after inoculation, showed evidence of discomfort, and extensive granulomatous inflammation at the inoculation sites. Lesions were minimal to mild in rabbits given antigen with RA and PBS with either adjuvant; While RA did not result in adverse side effects, the IgG response to (TG)-AL with RA was transient compared to FA.